

Quasiperiodic propagation in time of some classical/quantum systems: Nielsen's conserved quantity and Floquet properties.

P Kramer, Theoretische Physik U. Tuebingen, Germany,
 T Kramer, Institute for Theoretical Physics, U. Regensburg, Germany,
 V I Man'ko, Lebedev Physical Institute, Moscow, Russia.

October 19, 2008

Abstract.

We consider classical and quantum propagators for two different time intervals. If these propagators follow one another in a Fibonacci sequence we get a discrete quasiperiodic system. A theorem due to Nielsen provides a novel conserved quantity for this system. The Nielsen quantity controls the transition between commutative and non-commutative propagation in time. The quasiperiodically kicked oscillator moreover is dominated by quasiperiodic analogues of the Floquet theorem.

1 Introduction.

For the general description of time-dependent systems in quantum mechanics we refer to [4], [19], [8], [9], [3]. We consider the time evolution for a particular class of time-dependent classical/quantum systems which are periodic or quasiperiodic in time.

We restrict our attention to time evolutions, built piecewise from strings with classical symplectic propagators $g(t)$ in $2D$ phase space. These describe oscillators and dilatations, or positive and negative δ -kicks. Quantum propagators are constructed as unitary representations $S_{g(t)}$ of the classical linear canonical transformations for $g(t)$ from the symplectic group $Sp(2, R)$.

We present Nielsen's theorem on automorphisms of the free group F_2 with two generators and from it derive a conserved quantity in terms of the group commutator. We analyze Floquet properties of classical and quantum propagators. We show the presence/absence of Floquet quantum numbers for periodic system under the control

of system parameters. For the periodically kicked classical oscillator we demonstrate the occurrence of Floquet bands and Floquet gaps.

For periodic systems built from two basic intervals, we construct Nielsen's conserved quantity as a function of the system parameters. We extend the analysis to quasiperiodic strings. By building the strings through a Fibonacci automorphism we are able to survey their quasi-Floquet properties.

2 Floquet theorem for linear differential equations.

1 Theorem Floquet: Consider the real first-order periodic linear system of deq.

$$\frac{d}{dt}Y(t) = A(t)Y(t), A(t) \text{ real}, A(t+T) = A(t), T > 0. \quad (1)$$

A fundamental system of two solutions can be written in the form

$$Y(t) = \exp(\pm iQt)Y_0(t), Y_0(t+T) = Y_0(t), 0 \leq Q < \frac{2\pi}{T}. \quad (2)$$

We denote the real number Q as the Floquet index and $\exp(iQT)$ as the Floquet factor. It will become clear in what follows that the Floquet theorem requires complex solutions of the deq.1 for real $A(t)$.

Proof: The discrete time translation $T : A(t) \rightarrow A(t+T)$ is a symmetry operation of $A(t)$. It generates the abelian group C_∞ with elements $\pm nT, n = 0, 1, \dots$. The unitary representations of this group have the form

$$T \rightarrow \exp(i\Omega T), D^\Omega(nT) = \exp(in\Omega T), 0 \leq \Omega < \frac{2\pi}{T}. \quad (3)$$

The orthogonality and completeness relation of C_∞ are given in Appendix A. Consider a solution of eq. 2 which can be arranged so that it transforms according to an irreducible representations of C_∞ ,

$$Y^\Omega(t) : Y^\Omega(t+T) = Y^\Omega(t) \exp(i\Omega T). \quad (4)$$

From this solution define the function

$$Y_0^\Omega(t) := \exp(-i\Omega t)Y^\Omega(t) \quad (5)$$

Then it easily follows that $Y_0^\Omega(t)$ is periodic,

$$Y_0^\Omega(t+T) = Y_0^\Omega(t) \quad (6)$$

Since $A(t)$ is real, the complex conjugate solution $\overline{Y^\Omega} := Y^{-\Omega}$ is a second linearly independent solution with the complex conjugate Floquet factor. With $\Omega = Q$ we obtain eq. 2.

Remark (i): The Floquet theorem is mathematically equivalent to the Bloch theorem for systems with discrete position symmetry, which has the same abstract symmetry group C_∞ .

Remark (ii): We remark that the periodic system eq. 1 for certain ranges of its parameters cannot have solutions with the Floquet property. We shall give examples of this situation in later sections.

3 The free group F_2 and its automorphisms.

We refer to [18] for what follows. We also quote [7] as a useful reference to basic concepts for free monoids.

2 Def. The infinite *free group* F_2 consists of all words generated by concatenation from two non-commuting invertible generators $\{y_1, y_2\}$. Its elements can be viewed as a tree formed by words of increasing length.

3 Def: An automorphism of F_2 is an invertible homomorphic map

$$\phi : F_2 \rightarrow F_2, \{y_1, y_2\} \rightarrow \{\phi_1(y_1, y_2), \phi_2(y_1, y_2)\}, \{\phi_1, \phi_2\} \in F_2, \quad (7)$$

and so preserves group multiplication and unit element. The set of all automorphisms of F_2 forms the infinite group $\{\phi\} = \text{Aut}(F_2)$. This group, as was shown by Nielsen [18], is again finitely generated, with explicit generators given in [18]. Similar results hold true for the free group F_n and its group of automorphisms $\text{Aut}(F_n)$. Homomorphic properties of finitely generated groups have the virtue that they can be verified by checking them only for the generators.

Example 1: The Fibonacci automorphism is defined by

$$\begin{aligned} \phi_{fibo} : \{y_1, y_2\} &\rightarrow \{\phi_{fibo,1}(y_1, y_2), \phi_{fibo,2}(y_1, y_2)\} := \{y_2, y_1 y_2\}, \\ \phi_{fibo}^{-1} : \{y_1, y_2\} &\rightarrow \{y_2 y_1^{-1}, y_1\}, \end{aligned} \quad (8)$$

We shall use in particular iterated automorphisms as $(\phi_{fibo})^n$. Then the words $(\phi_{fibo})^n(y_1)$, with their length $|w|$, defined [7] as the number of letters in w after reducing w by use of all relations of the type $yy^{-1} = e$, become

$$\begin{array}{lcl} (\phi_{fibo})^n(y_1) : & y_1 & y_2 & y_1 y_2 & y_2 y_1 y_2 & y_1 y_2^2 y_1 y_2 & \dots \\ |(\phi_{fibo})^n(y_1)| : & 1 & 1 & 2 & 3 & 5 & \dots \end{array} \quad (9)$$

The word length $|(\phi_{fibo})^j(y_1)|$ is the Fibonacci number n_j

$$n_j : n_{j+2} = n_j + n_{j+1}, n_1 = n_2 = 1. \quad (10)$$

Notice the recursive structure of the images under $(\phi_{fibo})^n$: Define $(\phi_{fibo})^{(n-1)}(y_1) =: w_1$, $(\phi_{fibo})^{(n-1)}(y_2) =: w_2$. Then it follows from eq. 7 that $(\phi_{fibo})^n(y_1) = w_2$,

$(\phi_{fibo})^n(y_2) = (w_1 w_2)$, and so we must in step n simply concatenate the two words (w_1, w_2) obtained in step $(n - 1)$.

4 Nielsen theorem on $Aut(F_2)$: Consider the commutator $\mathcal{K} = y_1 y_2 y_1^{-1} y_2^{-1}$ in F_2 and a fixed automorphism $\phi(F_2) \in Aut(F_2)$. Then the image of the commutator under ϕ obeys, [18] theorem 3.9 p. 165,

$$\phi(\mathcal{K}) = w \mathcal{K}^{\pm 1} w^{-1}, \quad w \in F_2, \quad (11)$$

where the sign ± 1 and the element w depend on the chosen automorphism ϕ .

Proof: It suffices to prove the result for the finite set of generators of $Aut(F_2)$ given by Nielsen [18].

So the commutator $\mathcal{K}$ is conserved, up to inversion and conjugation, under any automorphism of $Aut(F_2)$. No counterpart of the Nielsen theorem 3 is known for $Aut(F_n), n > 3$.

Example 2: Evaluation of the image of the commutator under the particular Fibonacci automorphism ϕ_{fibo} yields

$$\mathcal{K} := y_1 y_2 y_1^{-1} y_2^{-1}, \quad \phi_{fibo}(\mathcal{K}) = y_2 (y_1 y_2) y_2^{-1} (y_1 y_2)^{-1} = y_2 y_1 y_2^{-1} y_1^{-1} = \mathcal{K}^{-1}. \quad (12)$$

The power of the Nielsen theorem appears through the notion of automorphisms induced on some group G .

5 Def: Induced automorphism: Consider a homomorphism $hom : F_2 \rightarrow G$ defined in terms of the generators by $hom : \{y_1, y_2\} \rightarrow \{g_1, g_2\} \in G$. The automorphism of G induced by the automorphism $\phi \in Aut(F_2)$ eq. 8 is the map $\phi_G : \{g_1, g_2\} \rightarrow \{\phi_1(g_1, g_2), \phi_2(g_1, g_2)\}$. Clearly induced automorphisms on G are homomorphic to the corresponding automorphisms from $Aut(F_2)$.

For an induced automorphism, the commutator induced in the group G has additional significance since it measures the non-commutativity of the group. In case $\mathcal{K} = I$ we have commutativity on the induced level which allows to rearrange the order of successive propagators.

Example 3: We shall consider in particular automorphisms induced on the symplectic group $Sp(2, R)$, the group of linear canonical transformations of a 1D classical hamiltonian system. For $g \in Sp(2, R)$, we have the algebraic equivalence relation $g^{-1} \sim g$ and the trace relation $\text{tr}(g^{-1}) = \text{tr}(g)$. Note that the character is a class function. If these properties are combined with the Nielsen theorem, one finds

6 Prop: For any automorphism of the free group F_2 induced on $Sp(2, R)$, the half-trace of the commutator,

$$\frac{1}{2} \text{tr}(\mathcal{K}) = \frac{1}{2} \text{tr}(g_1 g_2 g_1^{-1} g_2^{-1}) \quad (13)$$

is a quantity conserved under iterated automorphisms.

The condition $\frac{1}{2} \chi(\mathcal{K}) = 1$ is a necessary but not a sufficient condition for $\mathcal{K} = e$. So commutativity requires extra checking.

If by iterated automorphisms we can generate a string of arbitrary length, the conserved quantity eq. 13 can be assigned to the infinite string.

If we choose the Fibonacci automorphism eq. 8 and associate two matrix propagators to its two intervals, we generate a quasiperiodic system. We term such a system discrete quasiperiodic. Such system form a special class among general quasiperiodic systems [16].

In later sections, we shall associate to the classical system a quantum system. To a symplectic matrix g we shall associate a representation by a unitary operator $U(g)$ in Hilbert space.

Since this association is a homomorphic map from symplectic matrices into propagators which represent linear canonical transformations, the Nielsen theorem then extends to a theorem in Hilbert space.

4 Geometry and dynamics of systems of traces.

The traces of the symplectic matrices $g \in Sp(2, R)$ obviously play an important part in the classical and quantum analysis. For a subclass of systems, the computation of these traces does not even require the computation of the underlying matrices [14]. The Fibonacci system Example 1 belongs to this class. The trace invariant eq. 13 geometrically becomes a cubic surface in 3-space whose shape depends on the value of the invariant. The traces under repeated automorphisms and with increasing string length determine recursively a discrete dynamical system on this conserved surface. We sketch this approach: For the Fibonacci system eq. 8 we define $y_3 := y_1 y_2$ and $g_3 := g_1 g_2$. With the induced matrix automorphism written as

$$(X^n, Y^n, Z^n) := (\phi^n(g_1), \phi^n(g_2), \phi^n(g_3)) \quad (14)$$

and the matrix decomposition

$$g \in Sp(2, R) : g = S(g) + A(g), \quad S = \frac{1}{2} \text{trace}(g)e, \quad A = g - S, \quad \text{trace}(A) = 0 \quad (15)$$

one finds from eqs. (57-63) of [14] the recursion relations and the conserved quantity

$$\begin{bmatrix} S(X^{(n+1)}) \\ S(Y^{(n+1)}) \\ S(Z^{(n+1)}) \end{bmatrix} = \begin{bmatrix} S(Y^n) \\ S(Z^n) \\ 2S(Y^n)S(Z^n) - S(X^n) \end{bmatrix} \quad (16)$$

$$S(\mathcal{K}) = -4S(g_1)S(g_2)S(g_3) + 2[(S(g_1)S(g_1) + S(g_2)S(g_2) + S(g_3)S(g_3))] - e$$

All the quantities in eq. 16 by eq. 15 can be expressed in terms of real traces. The recursion relation in eq. 16 determines a discrete dynamical system with initial data given by the traces of (g_1, g_2, g_3) . By use of eq. 15, the conserved quantity $\frac{1}{2}\chi(\mathcal{K})$ can be expressed in terms of the three initial traces. We refer to [17] for a view of the cubic surface which represents $\frac{1}{2}\chi(\mathcal{K})$.

5 Quadratic hamiltonians and propagators.

The quantum propagators for quadratic hamiltonians are given in [10], [11]. They are unitary integral operators $S_{g(t_2-t_1)}$ where $g(t)$ is a linear canonical transformation $g \in Sp(2m, R)$, which can be determined by exponentiation from the action of the hamiltonian on the canonical position and momentum operators. So for quadratic hamiltonians we have a very clear correspondence between classical and quantum propagation: The classical time evolution is given by the action of $g(t)$ on the initial canonical coordinates at time zero. The quantum propagator is given by the action of $S_{g(t)}$ on the initial state at time zero.

6 Hamiltonians and propagators for $n = 1$.

In [10],[11] one finds for quadratic hamiltonians the matrix analysis of propagators in various dimensions m . The classification of types of hamiltonians for $n = 1, 2$ is discussed in detail in [21]. We briefly summarize the well-known results for $m = 1$. For $m = 1$, that is for 1-dimensional position space, we have $g(t) \in Sp(2, R)$. This group is particularly transparent. Its classes admit a subdivision into three types, depending on the value of their half-trace $\frac{1}{2}\chi(g) := \frac{1}{2}tr(g)$. Since $\det(g) = 1$, the eigenvalues of g are completely determined by the trace. Moreover the trace is independent of symplectic similarity transformations, $\chi(qgq^{-1}) = \chi(g)$, and so allows to determine the class of the symplectic matrix g . For $I : \frac{1}{2}|\chi(g)| < 1$, g is symplectically equivalent to the dynamics generated by an oscillator hamiltonian,

$$I : H = \frac{1}{2m}P^2 + \frac{m\omega}{2}X^2, \quad (17)$$

for $II : \frac{1}{2}|\chi(g)| > 1$ they are equivalent to an inverted oscillator or a dilatation

$$II : H = \kappa \frac{1}{2}(PX + XP) = \kappa(XP + \frac{h}{2i}), \quad (18)$$

and for $III : \frac{1}{2}|\chi(g)| = 1$ they are, apart from the extra case $g = e$, $\frac{1}{2}\chi(e) = 1$, equivalent to a free motion,

$$III : H = \frac{1}{2m}P^2, \quad (19)$$

All three hamiltonians are hermitian and time-independent. Hence their propagators $S_g(t)$ are unitary operators. For type I and III the propagators can be written as integral operators according to Moshinsky and Quesne [20] in the Schroedinger representation or in the Bargmann space of analytic functions, compare [10].

Type I: Harmonic oscillator.

Rewrite the harmonic oscillator hamiltonian eq. 17 as

$$H = \frac{1}{2}\omega \left[\frac{1}{m\omega} P^2 + m\omega X^2 \right]. \quad (20)$$

The corresponding linear canonical transformation is obtained as

$$\begin{aligned} g_\omega(t) &= \begin{bmatrix} \frac{1}{\sqrt{m\omega}} & 0 \\ 0 & \sqrt{m\omega} \end{bmatrix} \begin{bmatrix} \cos(\omega t) & \sin(\omega t) \\ -\sin(\omega t) & \cos(\omega t) \end{bmatrix} \begin{bmatrix} \sqrt{m\omega} & 0 \\ 0 & \frac{1}{\sqrt{m\omega}} \end{bmatrix}, \\ &= \begin{bmatrix} \cos(\omega t) & \frac{1}{m\omega} \sin(\omega t) \\ -(m\omega) \sin(\omega t) & \cos(\omega t) \end{bmatrix} \end{aligned} \quad (21)$$

Eq. 21 admits the complex diagonalization

$$\begin{aligned} &\begin{bmatrix} \cos(\omega t) & \sin(\omega t) \\ -\sin(\omega t) & \cos(\omega t) \end{bmatrix} \\ &= \sqrt{\frac{1}{2}} \begin{bmatrix} 1 & i \\ 1 & -i \end{bmatrix} \begin{bmatrix} \exp(-i\omega t) & 0 \\ 0 & \exp(i\omega t) \end{bmatrix} \sqrt{\frac{1}{2}} \begin{bmatrix} 1 & 1 \\ -i & i \end{bmatrix} \end{aligned} \quad (22)$$

In the complex domain we can therefore pass to a pair of classical variables which propagate with $\exp(\mp i\omega t)$.

The corresponding quantum propagator in the Bargmann space of analytic functions is [10]

$$\langle z' | S_{g_\omega(t)} | z \rangle = \exp(-i\omega t/2) \exp(\exp(i\omega t) z' \bar{z}). \quad (23)$$

The propagator in the Schroedinger representation is given in [10] pp. 34-6.

Type II: The hyperbolic and dilatation hamiltonian.

For type II the symplectic matrix may be taken as

$$g(t) = \begin{bmatrix} \exp(\kappa t) & 0 \\ 0 & \exp(-\kappa t) \end{bmatrix} \quad (24)$$

Note the correspondence to eq. 22 for $\kappa \rightarrow -i\omega$.

There is for type II no obvious integral operator representation. In this case we proceed directly from the hamiltonian eq. 18 and find

$$\exp(-\kappa t \frac{i}{h} H) \psi(x) = \exp(-\kappa \frac{t}{2}) (\exp(-\kappa t x \frac{d}{dx}) \psi)(x) = \exp(-\kappa \frac{t}{2}) \psi(\exp(-\kappa t) x). \quad (25)$$

This propagator is a time-dependent unitary dilatation. Applied to the ground state of an oscillator with frequency ω_0 , it transforms it into an oscillator ground state with frequency $\omega(t) = \omega_0 \exp(-2\kappa t)$. As a function of time, the uncertainty in position would increase while the uncertainty of momentum would decrease.

Another form for class *II* is given by the hyperbolic hamiltonian

$$H = \frac{1}{2} \left[\frac{P^2}{m} - m\kappa^2 X^2 \right]. \quad (26)$$

with the symplectic propagator

$$g_\kappa(t) = \begin{bmatrix} \cosh(\kappa t) & (m\kappa)^{-1} \sinh(\kappa t) \\ (m\kappa) \sinh(\kappa t) & \cosh(\kappa t) \end{bmatrix} \quad (27)$$

By choosing at time $t = 0$ the initial matrix

$$h(0) = \begin{bmatrix} 1 & 0 \\ 0 & m\kappa \end{bmatrix}, \det(h(0)) = m\kappa, \quad (28)$$

we determine two canonical column solutions given by

$$\begin{bmatrix} X_I(t) & X_{II}(t) \\ P_I(t) & P_{II}(t) \end{bmatrix} = g_\kappa(t)h(0) = \begin{bmatrix} \cosh(\kappa t) & \sinh(\kappa t) \\ (m\kappa) \sinh(\kappa t) & (m\kappa) \cosh(\kappa t) \end{bmatrix} \quad (29)$$

Inserting the fundamental two solutions into the hamiltonian eq. 26 we find the two different energies

$$E_I(0) = -\frac{m\kappa^2}{2}, \quad E_{II}(0) = \frac{m\kappa^2}{2}. \quad (30)$$

The types *I*, *II* of propagation in phase space are shown in Fig. 1.




Fig. 1. **Phase space representation.** The harmonic and the hyperbolic column solutions I: eq. 21 and II: eq. 27 in phase space.

In appendix B we give the relation between the propagator and the transfer matrix.
Type III: Free particle. From the hamiltonian

$$H = \frac{1}{2m} P^2 \quad (31)$$

one finds [10] the symplectic matrix

$$g(t) = \begin{bmatrix} 1 & -t \\ 0 & 1 \end{bmatrix} \quad (32)$$

and the propagator

$$\langle x' | S_{g(t)} | x \rangle = \exp(-i\pi/4) \sqrt{\frac{m}{2\pi\hbar t}} \exp(i \frac{(x' - x)^2}{2t}). \quad (33)$$

7 Prop: The types I, II, III together with the case $g = e$ exhaust, up to conjugation within $Sp(2, R)$, the symplectic propagators with quadratic hamiltonians.

7 Quadratic piecewise time-independent hamiltonians.

We consider now a system whose hamiltonian for $0 \leq t \leq T = T_1 + T_2$ is time-independent on the two intervals. Assume that the propagation for the full interval T is repeated periodically. Over the period we have then a propagation with the half-trace given by eq. 42.

For $\frac{1}{2}|\chi| < 1$ we can write $\frac{1}{2}\chi = \cos(\Omega T)$ and determine $\Omega, 0 \leq \Omega < 2\pi/T$. This value Ω is the Floquet index of a classical symplectic periodic system. The two Floquet factors $\exp(\pm i\Omega T)$ appear as the eigenvalues of the symplectic matrix $g_1(T_1)g_2(T_2)$ and determine the classical propagators $U(T)$. The Floquet index is completely determined by the parameters $(\omega_1, T_1, \omega_2, T_2)$ of the system. At multiples $nT, n = 1, 2, \dots$ the classical propagator becomes $(U(T))^n$.

In the quantum propagation, Ω represents the overall frequency of an effective oscillator hamiltonian. The propagator over the period T from eq. 23 becomes

$$\langle z' | S_{g_c(T)} | z \rangle = \exp(-i\Omega T/2) \exp(\exp(i\Omega T) z' \bar{z}), \quad (34)$$

We emphasize that this result holds true only for $T = T_1 + T_2$ whereas at intermediate time the propagator is governed by $g_1(t)$ or $g_2(t)$ respectively. The classical Floquet parameter enters the quantum propagator in an essential way. The eigenstates $|N\rangle$ of the overall harmonic oscillator of frequency Ω from eq. 34 are the monomials of degree N . These eigenstates under the discrete time translations transform according to

$$T : |N\rangle \rightarrow |N\rangle \exp(iN\Omega T) \quad (35)$$

This again generates an irreducible representation of the time translation group. By the unique decomposition $N\Omega = m2\pi/T + \Omega', 0 \leq \Omega' < 2\pi/T, m = 0, 1, 2, \dots$, this representation becomes $D^\Omega(NT) = D^{\Omega'}(T)$.

It follows that not the quantum propagator itself, but only the eigenstates in its decomposition transform irreducibly under the period T .

For $\frac{1}{2}|\chi| > 1$ the situation changes. By writing $\frac{1}{2}|\chi| = \cosh(\kappa T)$ we determine an effective classical dilatation parameter κ . The eigenvalues $\exp(\pm\kappa T)$ represent this dilatation over one period. The corresponding classical propagation does not represent a Floquet propagation since the eigenvalues of the symplectic matrix do not have absolute value 1. They form a non-unitary real representation of the time translation group. The n th powers of these two factors now approach $0, \infty$ respectively.

The unitary quantum propagator over a period T from eq. 25 becomes

$$\exp(-\kappa T \frac{i}{\hbar} H) \psi(x) = \exp(-\kappa \frac{T}{2}) (\exp(-\kappa T x \frac{d}{dx}) \psi)(x) = \exp(-\kappa \frac{T}{2}) \psi(\exp(-\kappa T)x). \quad (36)$$

Again, this propagator is valid only at time $T = T_1 + T_2$. For intermediate time the propagator is governed by one of the oscillator matrices $g_1(t)$ or $g_2(t)$ respectively. The quantum propagator over the periods T stays unitary, but for $n \rightarrow \infty$ when acting on a state transforms it up to normalization into a δ -distribution of sharp effective momentum/position.

Remark (i): The results on time-periodic systems from piecewise quadratic hamiltonians resemble the analysis of position-periodic Kronig-Penney systems [12] with piecewise constant potentials. In the latter systems, the potentials play the role of the frequencies, and the Bloch k -label plays the role of the Floquet index. Real values of the Bloch parameter yield a band index, while imaginary values indicate exponentially increasing or decreasing solutions of the Schroedinger equation which are excluded as band gaps.

Example 5: For the system of section 8 in case $|\frac{1}{2}\chi| < 1$ we obtain an overall Floquet state with Ω determined by $\frac{1}{2}\chi = \cos(\Omega T)$. Each Floquet state is the analogue to a Bloch state of a band model in x -space.

8 Oscillators and δ -kicks.

The transfer and propagator matrix methods apply to sequences of time intervals.

Example 6: The classical square well for the time interval t has the type I oscillator propagator

$$g_\omega(t) = \begin{bmatrix} \cos(\omega t) & (m\omega)^{-1} \sin(\omega t) \\ -(m\omega) \sin(\omega t) & \cos(\omega t) \end{bmatrix} \quad (37)$$

The initial values for the two fundamental column solutions I, II we take as

$$h(0) = \begin{bmatrix} 1 & 0 \\ 0 & m\omega \end{bmatrix} \quad (38)$$

Then the energies computed by insertion into the harmonic oscillator hamiltonian become

$$E_I = E_{II} = \frac{m\omega^2}{2}. \quad (39)$$

Example 7: The square tunnel for the time interval t has the repulsive oscillator type *II* propagator eq. 27. The energy of the two fundamental solutions are $E_I = -\frac{m\kappa^2}{2}$, $E_{II} = \frac{m\kappa^2}{2}$.

Following what is done in space in [13], [14] we introduce w.r.t. time, negative and positive δ -kicks. They are obtained as limits of square well and square tunnel intervals of finite length. In the limits : $m\omega^2 T \rightarrow u'$, $\omega T \rightarrow 0$ and $m\kappa^2 T \rightarrow u'$, $\kappa T \rightarrow 0$ with u the strength parameter we get from eqs. 37, 38 respectively the symplectic transfer matrices for negative/positive δ -kicks in the form

$$g_{\pm u'} = \begin{bmatrix} 1 & 0 \\ \mp u' & 1 \end{bmatrix}. \quad (40)$$

9 The periodically kicked oscillator.

We employ here the transfer matrix, see eq. 56, with $u' = mu$,

$$M(t) = \lambda_i = \begin{bmatrix} \cos(\omega T_i) & \omega^{-1} \sin(\omega T_i) \\ -\omega \sin(\omega T_i) & \cos(\omega T_i) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ u & 1 \end{bmatrix}, \quad i = 1, 2. \quad (41)$$

with positive δ -kicks. The half-trace becomes

$$\chi = \frac{1}{2} \text{Tr}(M(T)) = \cos(\omega T) + \frac{u}{2\omega} \sin(\omega T) \quad (42)$$

This expression has the period $\frac{2\pi}{\omega}$. We find

$$\chi = 0 : \cot(\omega T_0) = -\frac{u}{2\omega} \quad (43)$$

The solutions T_0 of this equation exist in each interval $2m\pi \leq \omega T \leq 2(m+1)\pi$, $m = 0, \pm 1, \pm 2, \dots$ and mark the centers of Floquet bands as functions of κ . The equations

$$\begin{aligned} \frac{1}{2}\chi = +1 : \tan\left(\frac{\omega T_+}{2}\right) &= \frac{u}{2\omega}, \\ \frac{1}{2}\chi = -1 : \cot\left(\frac{\omega T_-}{2}\right) &= -\frac{u}{2\omega}, \end{aligned} \quad (44)$$

determine for any such band its edges $T_{\pm}$ as functions of κ . Note that these equations have in $\omega T_{\pm}$ the period 4π . In between two Floquet bands there are gaps. In these gaps we have $\frac{1}{2}|\chi| > 1$ and so solutions of type II.

7 Prop: The Floquet spectrum of the periodically kicked oscillator. The periodically kicked oscillator has a Floquet band spectrum. Each interval $2m\pi \leq \omega T \leq 2(m+1)\pi$, $m = 0, 1, 2, \dots$ carries two bands. The centers T_0 from eq. 43 are repeated periodically, the edges given by eq. 44. The bands are separated by gaps which admit only solutions of type II.

Example 8: A simple example is provided by the choice $\frac{u}{2\omega} = 1$. Then the half-trace $\frac{1}{2}\chi$ eq. 42 with $\beta := \omega T$ becomes

$$\frac{1}{2}\chi = \cos(\beta) + \sin(\beta) = \sqrt{2} \cos(\beta - \pi/4). \quad (45)$$

The band edges β^+, β^- and the band centers β^0 are located at

$$\begin{aligned} \beta^+ &: 0 + 2\pi m_1, \quad \frac{\pi}{2} + 2\pi m_2, \\ \beta^- &: \pi + 2\pi m_3, \quad \frac{3\pi}{2} + 2\pi m_4, \\ \beta^0 &: \frac{\pi}{4} + \pi m_5, \quad m_j = 0, \pm 1, \pm 2, \dots \end{aligned} \quad (46)$$

Bands and gaps alternate with the same width $\frac{\pi}{2}$, as shown in Fig 2.




Fig. 2. **Floquet band structure for the periodically kicked oscillator.** Shown is the half-trace $\frac{1}{2}\chi$ eq. 42 as a function of ωT for the parameter $\frac{u}{2\omega} = 1$ of eq. 45. Bands and gaps alternate periodically and both have the width $\Delta\beta = \frac{\pi}{2}$.

10 The quasiperiodically kicked oscillator.

For properties of systems with a general quasiperiodic Hamiltonian we refer to [5]. The Nielsen theorem is valid for all discrete quasiperiodic systems of Fibonacci type, but its significance can already be seen on a simple example. Here we study a specific discrete quasiperiodic system of Fibonacci type in time, in analogy to a system in space from [12]. Following [6], it would be worthwhile to study the spectral function for this discrete quasiperiodic system.

For two intervals $(1, 2)$ which form a quasiperiodic Fibonacci sequence we take the matrix propagators in the form

$$\lambda_i = \begin{bmatrix} \cos(\omega T_i) & \omega^{-1} \sin(\omega T_i) \\ -\omega \sin(\omega T_i) & \cos(\omega T_i) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ u & 1 \end{bmatrix}, \quad i = 1, 2. \quad (47)$$

Then we compute, expanded in powers of $\frac{u}{\omega}$, the products

$$\begin{aligned} \lambda_1 \lambda_2 &= \begin{bmatrix} \cos(\omega(T_1 + T_2)) & \omega^{-1} \sin(\omega(T_1 + T_2)) \\ -\omega \sin(\omega(T_1 + T_2)) & \cos(\omega(T_1 + T_2)) \end{bmatrix} \\ &+ \frac{u}{\omega} \begin{bmatrix} 2 \sin(\omega T_1) \cos(\omega T_2) + \cos(\omega T_1) \sin(\omega T_2) & \omega^{-1} \sin(\omega T_1) \sin(\omega T_2) \\ \omega(2 \cos(\omega T_1) \cos(\omega T_2) - \sin(\omega T_1) \sin(\omega T_2)) & \cos(\omega T_1) \sin(\omega T_2) \end{bmatrix} \\ &+ \left(\frac{u}{\omega}\right)^2 \begin{bmatrix} \sin(\omega T_1) \sin(\omega T_2) & 0 \\ \omega \cos(\omega T_1) \sin(\omega T_2) & 0 \end{bmatrix} \\ (\lambda_2 \lambda_1)^{-1} &= \begin{bmatrix} \cos(\omega(T_1 + T_2)) & -\omega^{-1} \sin(\omega(T_1 + T_2)) \\ \omega \sin(\omega(T_1 + T_2)) & \cos(\omega(T_1 + T_2)) \end{bmatrix} \\ &+ \frac{u}{\omega} \begin{bmatrix} \sin(\omega T_1) \cos(\omega T_2) & -\omega^{-1} \sin(\omega T_1) \sin(\omega T_2) \\ -\omega(2 \cos(\omega T_1) \cos(\omega T_2) - \sin(\omega T_1) \sin(\omega T_2)) & 2 \cos(\omega T_1) \sin(\omega T_2) + \sin(\omega T_1) \cos(\omega T_2) \end{bmatrix} \\ &+ \left(\frac{u}{\omega}\right)^2 \begin{bmatrix} 0 & 0 \\ -\omega \sin(\omega T_1) \cos(\omega T_2) & \sin(\omega T_1) \sin(\omega T_2) \end{bmatrix} \end{aligned} \quad (48)$$

For the commutator $\mathcal{K}$ and its inverse we find

$$\begin{aligned} \mathcal{K} &= (\lambda_1 \lambda_2)(\lambda_2 \lambda_1)^{-1} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \frac{u}{\omega} \sin(\omega(T_1 - T_2)) \begin{bmatrix} \cos(\omega(T_1 + T_2)) & -\omega^{-1} \sin(\omega(T_1 + T_2)) \\ -\omega \sin(\omega(T_1 + T_2)) & -\cos(\omega(T_1 + T_2)) \end{bmatrix} \\ &+ \left(\frac{u}{\omega}\right)^2 \sin(\omega(T_1 - T_2)) \begin{bmatrix} \sin(\omega T_1) \cos(\omega T_2) & -\omega^{-1} \sin(\omega T_1) \sin(\omega T_2) \\ \omega \cos(\omega T_1) \cos(\omega T_2) & -\cos(\omega T_1) \sin(\omega T_2) \end{bmatrix}, \\ \mathcal{K}^{-1} &= (\lambda_2 \lambda_1)(\lambda_1 \lambda_2)^{-1} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \frac{u}{\omega} \sin(\omega(T_1 - T_2)) \begin{bmatrix} -\cos(\omega(T_1 + T_2)) & \omega^{-1} \sin(\omega(T_1 + T_2)) \\ \omega \sin(\omega(T_1 + T_2)) & \cos(\omega(T_1 + T_2)) \end{bmatrix} \\ &+ \left(\frac{u}{\omega}\right)^2 \sin(\omega(T_1 - T_2)) \begin{bmatrix} -\cos(\omega T_1) \sin(\omega T_2) & \omega^{-1} \sin(\omega T_1) \sin(\omega T_2) \\ -\omega \cos(\omega T_1) \cos(\omega T_2) & \sin(\omega T_1) \cos(\omega T_2) \end{bmatrix}, \end{aligned} \quad (49)$$

Remember that in each iteration of the Fibonacci automorphism the commutator $\mathcal{K}$ by eq. 12 is transformed into its inverse. This does not affect the trace which becomes an invariant under the automorphism.

The half-trace of the commutator therefore becomes the invariant

$$\mathcal{I} = \frac{1}{2}\text{Tr}(\mathcal{K}) = \frac{1}{2}\text{Tr}(\mathcal{K}^{-1}) = 1 + \frac{1}{2}\left(\frac{u}{\omega}\right)^2(\sin(\omega(T_1 - T_2)))^2. \quad (50)$$

Of interest are the points where the invariant takes the value $I = 1$. These are given from eq. 50 by

$$\begin{aligned} \sin(\omega(T_1 - T_2)) &= 0, \\ \omega(T_1 - T_2) &= m\pi, m = 0, \pm 1, \pm 2, \dots \end{aligned} \quad (51)$$

For the Fibonacci case $T_1 = T, T_2 = \tau T$ we find from eqs. 50, 51

$$\omega(\tau - 1)T = m\pi, m = 0, \pm 1, \pm 2, \dots \quad (52)$$

The full commutator $\mathcal{K}$ at these points from eq. 49 becomes

$$\mathcal{K} = I. \quad (53)$$

and we find

8 Theorem: The invariant $\mathcal{I}$ as a function of T_1, T_2 has the value $\mathcal{I} = 1$ at the points eqs. 51 and 52. These points are repeated periodically. At each of these points, the commutator $\mathcal{K}$ becomes the unit matrix and so the two matrices eq. 47 commute.

For the Fibonacci model of eq. 52 we get periodic points where the two matrices λ_1, λ_2 commute. This does not automatically imply that we have a quasiperiodic Floquet propagation. To assure that, we must for both matrices be in an individual Floquet band. The relevant parameters for the individual Floquet bands are $\omega T, \tau\omega T$ respectively which scale by τ . To find overall propagation we must superimpose the two band structures and find the regions of band overlap. Moreover for commutativity we must fulfill the condition eq. 52. The superposition of the bands and the location of commutativity are shown in Fig 3.




Fig. 3. **Quasiperiodic Fibonacci system.** Bottom: For the parameters of eq. 46, the half-traces of the two systems exhibit two systems of bands as functions of ωT . Top: Overlap regions of the two systems of bands and the invariant $\mathcal{I}(\omega T)$. If a commutative point with $\mathcal{I} = 1$ hits a band overlap region, the system shows quasiperiodic Floquet behaviour. These points are marked by vertical arrows.

The propagation in phase space for multiple values of a commutative in-band case is shown in Fig. 4. All the points reached fall on a single ellipse equivalent to a harmonic oscillator.




Fig. 4. Quasiperiodic Floquet theorem: For multiples of a value ωT with $\mathcal{I}(\omega T) = 1$, hence commutative, and moreover in the overlap of the two Floquet bands, all phase space propagators fall on an ellipse equivalent to a harmonic oscillator.

11 Conclusion.

The algebraic Nielsen theorem applied to classical and quantum quasiperiodic systems yields a novel conserved quantity. This quantity as a continuous function of system parameters controls transitions between commutative and non-commutative propagation in time. The propagation of the quasiperiodically kicked oscillator demonstrates an interplay between the Nielsen and Floquet properties.

Acknowledgement.

One of the authors (P.K.) thanks Ch. Fulton, Dept. of Mathematical Sciences, Florida Institute of Technology, Melbourne, Florida, for helpful comments. Fulton pointed out reference [5] for quasiperiodic Hamiltonians, and the need to study the spectral function for quasiperiodic Sturm-Liouville problems.

Appendix A: Irreps of the discrete translation group.

We give here the orthogonality and completeness relations for the discrete translation group C_∞ as discussed in [15]. In the present case they apply to Floquet states. The orthogonality relation is

$$\begin{aligned} D^\Omega(nT) &= \exp(2\pi i \Omega nT), \\ \Omega &\in BZ = \{0, 2\pi/T\}, \quad n \in \{0, \pm 1, \pm 2, \dots\} \\ \sum_n \overline{D^\Omega(nT)} D^{\Omega'}(nT) d\Omega &= |BZ| \delta(\Omega - \Omega'). \end{aligned} \tag{54}$$

Eq. 54 yields an orthogonality relation between evolutions belonging to two different Floquet indices. It does not imply an orthogonality between pairs with the same Floquet index from different Floquet bands, as they are found for example in section 10. The completeness relation is

$$\int_{\Omega \in BZ} \overline{D^\Omega(nT)} D^\Omega(mT) d\Omega = |BZ| \delta_{n,m}. \tag{55}$$

Appendix B: Classical propagator and transfer matrix.

The matrix which propagates the canonical pair of position and momentum $\{X(t), P(t)\}$ we denote as the classical (symplectic) propagator $g(t) \in Sp(2, R)$, $\det(g(t)) = 1$. The matrix which propagates a fundamental pair of position and velocity $\{X(t), \dot{X}(t)\}$ we denote as the transfer matrix $M(t)$. Again we may choose $\det(M(t)) = 1$. For a hamiltonian with standard kinetic energy we have the relation $P(t) = m\dot{X}(t)$. From this we get the conjugation relation

$$M(t) = \begin{bmatrix} \sqrt{m} & 0 \\ 0 & \frac{1}{\sqrt{m}} \end{bmatrix} g(t) \begin{bmatrix} \frac{1}{\sqrt{m}} & 0 \\ 0 & \sqrt{m} \end{bmatrix} \tag{56}$$

between the symplectic propagator $g(t)$ and the transfer matrix $M(t)$. From the relation eq. 56 it follows that for a sequence of strings both their symplectic propagators and their transfer matrices are multiplied in the same order.

The first and second column of the propagator matrix $g(t)$ each yield a fundamental canonical pair $\{I, II\}$ of solutions and trajectories for the hamiltonian equations of motion. For quadratic hamiltonians, the hamiltonian equations of motion are linear. Then the most general classical solution is a linear superposition of fundamental solutions. The superposition changes the classical energy which therefore must be recalculated for any superposition.

References

- [1] Arnold V I, *Mathematical Methods of classical mechanics*, Springer, Berlin (1978)
- [2] Bargmann V, *Group representations in Hilbert spaces of analytic functions* (1968)
- [3] Bayfield, J E, *Quantum evolution*, Wiley, New York (1999)
- [4] Dodonov V V and Man'ko V I, Invariants and evolution of non-stationary quantum systems, ed. Markov, Nova Science Publishers, Cormack, N.Y. (1989)
- [5] Dinaburg E and Sinai Ya, *The One-dimensional Schrödinger Equation with a Quasiperiodic Potential*, Functional Anal. Appl, **9** (1975) 279-290
- [6] Fulton Ch, Pearson D, and Pruess, St, *Computing the spectral function for singular Sturm-Liouville problems*, J Comp. Appl. Math. **176** (2005) 131-162
- [7] Lothaire M, *Combinatorics on words*, Addison-Wesley, Reading (1983)
- [8] Karner G, Man'ko V I, and Streit L, Quasi-energies, loss-energies and stochasticity, Reports on Mathematical Physics, **29** (1991) 177-93
- [9] Karner G, Man'ko V I, and Streit L, Proc. Lebedev Physical Institute Nauka **208** (1992) 226
- [10] Kramer T, *Quantum ballistic motion in uniform electric and magnetic fields*, Diploma thesis, TU Munich (2000)
- [11] Kramer T, *Matter waves from localized sources in homogeneous force fields*, PhD thesis, TU Munich, 2003. Online: <http://tumb1.biblio.tu-muenchen.de/publ/diss/ph/2003/kramer.html>
- [12] Kramer P and Kramer T, *Exact electron states in 1D (quasi-) periodic arrays of δ -potentials*, in: *From quasicrystals to more complex systems* eds. F Axel et al, Springer, Berlin 2000, pp. 85-114.
- [13] Kramer P, *Energy gauge and electron confinement in quasicrystals*. J Phys **A** **31** (1998) 743-56
- [14] Kramer P, *Algebraic structure for one-dimensional quasiperiodic systems* J Phys **A** **26** (1993) 213-28
- [15] Kramer P and Lorente M, *Discrete and continuous symmetry via induction and duality*, in: *Symmetries in Science* **X**, B Gruber and M Ramek eds., Plenum, New York 1998, pp. 165-77

- [16] Kramer P, *Quasiperiodic systems*, in: Encyclopedia of Mathematical Physics, eds. J P Francoise, G L Naber, Sh Tsun Tsou, Elsevier, Amsterdam 2006, 308-315
- [17] Kohmoto M, Kadanoff L P and Tang C, Phys. Rev. Lett. **50** (1983) 1870
- [18] W Magnus, A Karrass, D Solitar, *Combinatorial group theory*, Dover, New York 1976
- [19] Malkin I A and Man'ko V I, *Dynamical symmetries and coherent states of quantum systems*, Nauka, Moscow (1979) (in Russian)
- [20] Moshinsky M and Quesne C, *Linear canonical transformations and their unitary representation*, J Math Phys **12** (1971) 1772-
- [21] Winternitz P and Moshinsky M, *Quadratic hamiltonians in phase space and their eigenstates*, J Math Phys **21** (1980) 1667-82